

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 47/48, 49/02		A2	(11) International Publication Number: WO 92/08494
			(43) International Publication Date: 29 May 1992 (29.05.92)
(21) International Application Number: PCT/US91/08253		(74) Agent: DAVIS, William, J.; 343 State Street, Rochester, NY 14650-2201 (US).	
(22) International Filing Date: 7 November 1991 (07.11.91)			
(30) Priority data: 610,861 8 November 1990 (08.11.90) US 784,333 29 October 1991 (29.10.91) US		(81) Designated States: AT (European patent), AU, BE (European patent), BR, CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), SU ⁺ .	
(71) Applicant: STERLING WINTHROP INC. [US/US]; 90 Park Avenue, New York, NY 10016 (US).		Published Without international search report and to be republished upon receipt of that report.	
(72) Inventors: TONER, John, L. ; 109 Brookhollow Drive, Downingtown, PA 19355 (US). HILBORN, David, A. ; 10 Southern Hill Circle, Henrietta, NY 14467 (US). MURRAY, Bruce, J. ; 1575 Sage Terrace, Walworth, NY 14568 (US). HOSSAIN, Timothy, Z. ; 13588 North Central Expressway, Dallas, TX 75243 (US).			

(54) Title: TARGETING RADIOACTIVE IMMUNOREAGENTS

(57) Abstract

A targeting radioactive immunoreagent comprises a metal radionuclide ion, a complexing agent, and an immunoreactive group covalently bonded to said complexing agent, the complexing agent having structure (A-I), wherein R represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group; R¹ represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group; R² represents hydroxy, carboxy, hydroxy-alkyl, carbonyliminodiacetic acid, methyleneiminodiacetic acid, methylenethioethyleneiminodiacetic acid, hydrazinylidenediacetic acid, or a salt of such acids, or two R² groups, taken together, represent the atoms necessary to complete a macrocyclic ring structure containing at least one heteroatom coordinating site and at least one, preferably two alkylene groups forming part of the ring structure; R³ represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group; R⁴ represents hydrogen or a protein reactive group; n is 0 to 4, o is 0 or 1, m is 0 or 1; provided that at least one of n and m is 0 and at least one of R, R¹, R³ and R⁴ is a protein reactive group. In a composition of matter aspect, this invention provides novel terpyridines and phenanthrolines. Preferred terpyridines have the structure (A-I) above wherein n = 1, m = 0 and R is a substituted phenyl containing an alkyl or alkoxy substituent and a protein reactive group. Preferred phenanthrolines have the structure (A-I) above wherein n = 0, m = 1 and at least one R⁴ is a protein reactive group. The targeting radioactive immunoreagents are particularly useful in therapeutic and diagnostic imaging compositions and methods.



+ DESIGNATIONS OF "SU"

Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU+	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

5

TARGETING RADIOACTIVE IMMUNOREAGENTS

10

FIELD OF THE INVENTION

15

The present invention relates to novel immunoreagents and more particularly to targeting radioactive immunoreagents which find particular utility in therapeutic and diagnostic imaging compositions and methods. The present invention further relates to novel complexing agents.

BACKGROUND OF THE INVENTION

20

25

30

Prior to 1980, the targeting of tumor-bearing sites by radioimmunoglobulin had been demonstrated by a number of laboratories at different institutions (S.E. Order et al., "Use of Isotopic Immunoglobulin in Therapy," Cancer Research 40, 3001-7 (August 1980)). By 1980 it was demonstrated that tumors would concentrate radiolabeled-antibodies to tumor associated antigens and that radiolabeled reagents employed allowed both diagnostic imaging of tumors, e.g., by gamma camera imaging (radioimmunoscinigraphy) and positron tomography, and therapeutic treatment, i.e., reduction in tumor size by the targeting radioactive immunoreagent.

35

Early targeting with radiolabeled immunoreagents was carried out with radioactive-iodine. However, as noted by Scheinberg et al, "Tumor Imaging with Radioactive Metal Chelates Conjugated to Monoclonal Antibodies," Science 215, No. 19, 1511-13 (March 1982), iodine isotopes pose several problems, particularly with respect to scanning of tumor images.

Of the three commonly available isotopic forms, only ^{123}I has the appropriate emission characteristics for imaging and a short enough half-life to be safely used diagnostically. The gamma radiation of ^{125}I is too weak for imaging. ^{131}I has often been used but is undesirable because of its long half-life and high energy gamma and cytotoxic beta radiations. ^{131}I has also been used therapeutically for large tumors, but appears ineffective in the treatment of small tumors. Moreover, rapid metabolism of radioiodinated antibodies allows incorporation of the iodine into the thyroid and active excretion of the iodine by the stomach and urinary tract. This dispersion of the radioactive iodine hinders imaging of specific tumors since the tumors are hidden by background radiation.

In addition to tumor targeting with radioactive antibodies for diagnostic imaging and therapeutic treatment, similar targeting has been accomplished for diagnostic imaging of infarcts, specifically, myocardial infarcts, using antibodies to canine cardiac myosin (Khaw et al, "Myocardial Infarct Imaging of Antibodies to Canine Cardiac Myosin Indium - 111 - Diethylenetriamine Pentaacetic Acid," Science 209, 295-7 (July 1980), and for imaging atherosclerosis by targeting atherosclerotic plaques. The same disadvantages in the use of radioactive iodine exist for diagnostic infarct imaging as for tumor imaging and therapeutic treatment.

It is known that ^{111}In can be complexed with polyaminocarboxylic acids such as ethylene diaminetetraacetic acid (EDTA) and diethylenetriaminepentaacetic acid (DTPA). However, the covalent linkage of proteins (antibodies) to these complexing agents, accomplished by acylation with activated carbonyls, aromatic diazonium coupling, or bromoacetylation is inefficient, even though the

isocyanatobenzyl derivatives described by Brechbiel et al "Synthesis of 1-(p-Isothiocyanatobenzyl)Derivatives of DTPA and EDTA. Antibody Labeling and Tumor Imaging Studies," Inorg. Chem. 25, 2772-81 (1986)) were created to facilitate covalent attachment of proteins with the complexing agents.

Recently, research efforts have been directed to improved antibodies (Ab's), e.g., monoclonal, specific antibodies for specific targeting, antibodies that complex or bind directly with radionuclides, preferred radionuclides and combinations thereof with antibodies and complexing agents. Some attempts have been made towards improving complexing agents.

Nonetheless, EDTA and especially DTPA and derivatives thereof have remained the prevalent complexing agents to covalently bind antibody and coordinately complex metallic radionuclides. However, the inadequacies of DTPA have been noted, for example, by Parker et al, "Implementation of Macrocyclic Conjugated Antibodies for Tumor Targeting," Pure and Appl. Chem., 61, No. 9, 1637-41 (1989)... "Conventionally the metal radionuclide has been complexed by an acyclic chelate (e.g. EDTA or DTPA) which is covalently linked to the antibody. None of the chelates is adequate because the metal tends to dissociate in vivo,"...and by Cox et al, "Synthesis of a Kinetically Stable Yttrium-90 Labelled Macrocyclic-Antibody Conjugate," J. Chem. Soc., Chem. Commun. 797-8 (1989)... "Yttrium-90 is an attractive isotope for therapy...but its clinical use will be very limited because of bone marrow toxicity, resulting from acid-promoted release of ⁹⁰Y from an antibody linked chelate such as diethylenetriamine-pentaacetic (DTPA)."

The attempts to develop improved complexing agents have provided materials which have their

shortcomings. For example, Craig et al "Towards Tumor Imaging with Indium-111 Labelled Macrocycle-Antibody Conjugates," J. Chem. Soc. Chem. Commun., 794-6 (1989) describe macrocyclic hexacoordinating ligands but state
5 that "The limiting feature of this approach is that ¹¹¹In labelling of the macrocycle is required before antibody conjugation. Indium binding by (4) is insufficiently fast at 37°C for efficient radiolabeling...Other tribasic triazamacrocyclic
10 ligands were screened therefore for their ability to bind indium rapidly under mild conditions (20°C, pH 5, < 1h), yet still form a kinetically stable complex in vivo...However, only (6) proved effective when the ligand concentration was 10μM, and under these
15 conditions a 96% radiolabeling yield was determined (30 min, pH 5, 20°C)."

Nevertheless, thirty minutes is still unsatisfactory. It would be highly desirable to have complexing agents superior to EDTA and DTPA which would
20 coordinately bind preferred radionuclides such as In, Y, Sc, Ga, Ge, etc. within a few minutes, i.e., in less than about 5 min, immediately prior to administration of the reagent to the patient, especially when a short-lived radionuclide must necessarily be generated from a
25 longer-lived radionuclide at the time of treatment of the patient.

It should be noted that complexes of yttrium, a preferred radionuclide, tend to be less stable than those of indium (Mather et al, "Labelling Monoclonal
30 Antibodies with Yttrium 90," Eur. J. Nucl. Med. 15, 307-312 (1989)) with respect to conventional complexes. Mather et al teach that biodistribution studies in cancer patients using radiolabeled antibodies have suggested that the in vivo stability of yttrium-labeled
35 antibodies is not as great as their ¹¹¹In-labelled counterparts and that these findings are supported by

other recent publications in the field.

When chelating agents are covalently bonded to proteins (Ab's), the proteins usually are capable of accepting far more than one molecule of the chelating agent because they contain a host of amine and sulfhydryl groups through which the chelating agents are bound. It is often very important to determine how many chelating sites are bound to each protein molecule. The most convenient way to accomplish this is by spectrophotometric means. However, prior art chelating agents and chelates thereof have spectra that overlap with those of useful proteins, and an analytical determination of the number of chelating or chelated sites per molecule of protein cannot be made by spectroscopy since the overlapping spectra mask each other. It would thus be highly desirable to obtain chelating agents for conjugation to proteins whose spectra, and whose metal chelate spectra, do not overlap with that of the proteins to which the chelating agents are chemically bonded.

Another problem with some prior art compositions is that the chelator must be activated by a reducing agent before forming the radionuclide chelate. If the protein conjugates are to be formed prior to formation of the radionuclide chelate, then the reducing agent employed for activating the complexing agent can degrade the protein. For example, the preferred chelating agents currently used for complexing technetium (Tc) and rhenium (Re) complex to the metals via sulfur-containing groups which must be reduced with a reducing agent (dithiothreitol) to activate the chelator before forming the radionuclide chelate. If the protein conjugate containing disulfide bonds is formed prior to reduction, then the reducing agent can degrade the protein. It would be highly desirable to have chelating agents capable of forming

conjugates with proteins before complexing with radionuclides, and particularly chelating agents for Tc and Re which do not require an activation step involving a reducing agent prior to complexation.

5 In summary, the various currently available radiolabeled antibodies and chelating agents employed for making immunoreactive conjugates by covalently bonding of a chelating agent to the immunoreactive protein, and radionuclide complexes thereof for use in
10 diagnostic imaging and targeted therapeutics suffer from certain of the following disadvantages:
1) toxicity; 2) dispersion of the reagent due to rapid metabolism; 3) inadequate emission characteristics; 4) inefficient covalent bonding with protein for conjugate
15 preparation; 5) slow complexation with metals; 6) unstable metal complexation, e.g., with respect to temperature, time or pH; 7) inability to form conjugates and store until metal complexation is desired; 8) inability to spectrophotometrically analyze
20 the radionuclide complex reagent; and 9) inability to complex without activation steps that degrade protein.


SUMMARY OF THE INVENTION

We have discovered targeting radioactive immunoreagents which solve the problems of the prior
25 art discussed above. The targeting radioactive immunoreagents of this invention comprise a metal radionuclide ion, a complexing agent which is a derivative of a pyridine, bipyridine, terpyridine, quaterpyridine, quinquepyridine, sexipyridine or
30 phenanthroline, and an immunoreactive group covalently bonded through a protein reactive group to the complexing agent.

More particularly, in accordance with the invention, there is provided:

35 a targeting radioactive immunoreagent comprising a metal radionuclide ion, a complexing

agent, and an immunoreactive group covalently bonded to the complexing agent, the complexing agent having the structure



wherein

R represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group;

R^1 represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group;

R^2 represents hydroxy, carboxy, hydroxyalkyl, carbonyliminodiacetic acid, methyleneiminodiacetic acid, methylenethioethyleneiminodiacetic acid, hydrazinylidenediacetic acid, or a salt of such acids, or two R^2 groups, taken together, represent the atoms necessary to complete a macrocyclic ring structure containing at least one heteroatom coordinating site and at least one, preferably two alkylene groups forming part of the ring structure;

R^3 represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy,

heterocyclyl or a protein reactive group;

R^4 represents hydrogen or a protein reactive group;

n is 0 to 4


o is 0 or 1

m is 0 or 1;

provided that at least one of n and m is 0 and at least one of R , R^1 , R^3 and R^4 is a protein reactive group.

The pyridines have the structure


A-II



wherein R^1 , R^2 and R^3 are as defined above.

The bipyridines, terpyridines, quaterpyridines, quinquepyridines and sexipyridines have the structure

A-III




wherein R , R^1 , R^2 and R^3 are as defined above and n is 0, 1, 2, 3 or 4.

The phenanthrolines have the structure


A-IV

A-IV



wherein R^2 , R^3 and R^4 are as defined above.

This invention provides novel terpyridines, preferably having the structure A-III above wherein $n=1$, R is



, R^5 is alkoxy or alkyl, and

R^6 is a protein reactive group.

This invention further provides novel phenanthrolines preferably having the structure A-IV above wherein at least one R^4 is a protein reactive group.

This invention also provides therapeutic and diagnostic compositions comprising the above-described targeting radioactive immunoreagent.

This invention further provides a method for diagnostic imaging a site in a patient comprising a) administering to the patient an effective amount of the above-described radioactive immunoreagent capable of targeting the site, and b) imagewise activating a radiation-sensitive element or device, such as, for example, a film or electronic sensor, with the radiation emitted from the targeted site.

A method for treating disease sites in a patient according to this invention comprises administering to the patient or a specimen from the patient an effective amount of a therapeutic composition comprising the above-described radioactive

immunoreagent capable of targeting the site and a pharmaceutically acceptable carrier therefor.

It is an advantageous feature of this invention that the described targeting radioactive immunoreagents exhibit lower toxicity, e.g., compared to other radioactive yttrium chelators.

It is an advantageous feature that the targeting immunoreagents of this invention are not rapidly metabolized and do not deleteriously disperse.

It is another advantageous feature that the described complexes efficiently form covalent bonds with proteins and other biological molecules.

Yet another advantageous feature of this invention is that the described immunoreagents exhibit good emission characteristics and are readily subject to spectrophotometric analysis.

Additionally, protein conjugates of the complexing agents can be formed and stored until metal complexation is desired and complexation can be accomplished without activation steps that degrade protein.

Moreover, the complexing agents rapidly complex with metals, and the resulting chelates exhibit excellent stability with respect to time, temperature and pH.

Other advantageous features of this invention will become readily apparent upon reference to the following description of the preferred embodiments, when read in light of the accompanying drawings.

Brief Summary of the Drawings

Fig. 1 depicts immunocompetency assays of B72.3-THT-Sc⁺⁺⁺, a radioactive immunoreagent of this invention, a B72.3-THT conjugate and unmodified B72.3.

Fig. 2 depicts immunocompetency assays of a B72.3-THT conjugate, two preparations of B72.3-TMT conjugates and unmodified B72.3.

Fig. 3 depicts the results of a biodistribution study of B72.3-TMT-¹¹¹In, a radioactive immunoreagent of the invention, and B72.3-DTPA-¹¹¹In.

5 Fig. 4 depicts the results of a biodistribution study of B72.3-TMT-⁹⁰Y, a radioactive immunoreagent of the invention, and B72.3-DTPA-⁹⁰Y.

Fig. 5 is the survival curve, i.e., the number of mice surviving each day past the first day of inoculation for a low dose regimen and a high dose
10 regimen.

Figs. 6 and 7 are spectra of TMT, TMT-Y⁺⁺⁺ and TMT-Pb⁺⁺.

Description of Preferred Embodiments

The description which follows primarily
15 concerns usage of the targeting radioactive immunoreagents in therapeutic and diagnostic imaging compositions and methods. In addition, the targeting radioactive immunoreagents are useful as diagnostic reagents, for example, radioimmuno-electrophoresis
20 reagents.

The immunoreagents of this invention comprise a metal radionuclide ion, a complexing agent, and an immunoreactive group covalently bonded to the complexing agent through a protein reactive group.

25 The complexing agent is a derivative of a pyridine, bipyridine, terpyridine, quaterpyridine, quinquepyridine, sexipyridine or phenanthroline, preferably having the structural formula A-I recited in the Summary above.

30 Each R in formula A-I independently is hydrogen; straight or branched alkyl, preferably containing from 1 to about 20 carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, s-butyl, t-butyl, 2-ethylhexyl, decyl, hexadecyl, octadecyl, etc.;
35 alkoxy, the alkyl portion of which contains from 1 to about 20 carbon atoms as described for R above;

alkylthio, the alkyl portion of which contains from 1 to about 20 carbon atoms as described for R above; alkylamino, the alkyl portion of which contains from 1 to about 20 carbon atoms as described for R above; 5 alkylformamido, the alkyl portion of which contains from 1 to about 20 carbon atoms as described for R above; substituted or unsubstituted aryl, preferably containing from about 6 to 20 carbon atoms such as phenyl, naphthyl, phenanthryl, nitrophenyl, 10 hydroxyphenyl, aminophenyl, hexadecylaminophenyl, octadecylaminophenyl, tolyl, xylyl, methoxyphenyl, 3-amino-4-methoxyphenyl, 4-methoxy-3-(N-methylhydrazinothioformamido)phenyl, 3-isocyanato-4-methoxyphenyl, 15 3-isothiocyano-4-methoxyphenyl, methylthiophenyl, carboxyphenyl and alkylaryl such as alkylphenyl, the alkyl portion of which contains from 1 to about 20 carbon atoms as described for R above; aryloxy, the aryl portion of which contains from 6 to 20 about 20 carbon atoms as described for R above; a substituted or unsubstituted heterocyclyl, preferably containing 5 to 6 nuclear carbon and heteroatoms such as N, S, P or O, such as pyridyl, methylpyridyl, 20 nitrophenyl, methoxypyridyl, oxazolyl, imidazolyl, pyrazolyl and quinolyl; or a protein reactive group. In especially preferred embodiments, R is a 4-alkoxy-3-aminophenyl or a 4-alkoxy-3-isothiocyano phenyl group.

Each R¹ independently is selected from the groups specified for R. R¹ preferably represents 30 hydrogen or a protein reactive group.

Each R² is independently selected from hydroxy; carboxy; hydroxyalkyl, the alkyl portion of which preferably contains from 1 to 4 carbon atoms, 35 such as hydroxymethyl;

carbonyliminodiacetic acid $[-\text{CON}(\text{CH}_2\text{COOH})_2]$;

methyleneiminodiacetic acid $[-\text{CH}_2\text{N}(\text{CH}_2\text{COOH})_2]$;

methylenethioethyleneiminodiacetic acid
 $[-\text{CH}_2\text{SCH}_2\text{CH}_2\text{N}(\text{CH}_2\text{COOH})_2]$;

- 5 hydrazinylylidenediacyetic acid, such as 1-hydrazinyl-2-ylidenediacyetic acid $[-\text{NHN}(\text{CH}_2\text{COOH})_2]$ and 1-methyl-1-hydrazinyl-2-ylidenediacyetic acid $[-\text{N}(\text{CH}_3)\text{N}(\text{CH}_2\text{COOH})_2]$;

- and 2,6-dicarboxy piperidino or the salts of such acids, including, for example, metal salts of such
10 acids formed from such metals as Na, K, Li, etc., and ammonium salts such as ammonium, tetraethylammonium, and tetramethylammonium salts. Alternatively, the two R^2 groups, taken together, represent the atoms necessary to complete a macrocyclic ring structure
15 containing (a) at least one heteroatom coordinating site for ions, and (b) at least one, preferably two alkylene groups forming part of the ring structure. The macrocyclic ring-forming groups can be a heteroatom group substituted alkylene such as 2,2-
20 bis(ethoxycarbonyl)-1,3-propylene; or heteroatom-containing groups such as oxybis(alkylene) such as oxybis(ethylene), oxybis(ethyleneoxymethylene), oxybis(ethyleneoxyethylene);
alkyleneoxyalkyleneoxyalkylene, such as
25 methyleneoxyethyleneoxymethylene; arylene-di(oxyalkylene), such as 1,4-dimethyl-5,6-phenylenebis(oxymethylene);

2,6-pyridylenebis(methyleneoxymethylene);

2-methoxy-5-methyl-1,3-phenylenebis(methyleneoxymethylene) and 1,10-phenanthroline-2,9-ylenebis(methyleneoxymethylene);

5 carboxymethyliminobis(trimethylenecarboxymethyliminomethylene) $[-CH_2N(CH_2COOH)(CH_2)_3N(CH_2COOH)-(CH_2)_3N(CH_2COOH)CH_2-]$;

carboxymethylthioethyliminobis(trimethylenecarboxymethylthioethyliminomethylene) $[-CH_2N(CH_2CH_2SCH_2COOH)_2-(CH_2)_3N(CH_2CH_2SCH_2COOH)_2-(CH_2)_3N(CH_2CH_2SCH_2COOH)CH_2-]$;

carboxymethyliminobis(ethylenecarboxymethyliminomethylene) $[-CH_2N(CH_2COOH)CH_2CH_2N(CH_2COOH)CH_2CH_2N(CH_2COOH)CH_2-]$; carboxymethylthioethyliminobis-

15 (ethylenecarboxymethylthioethyliminomethylene) $[-CH_2N(CH_2CH_2SCH_2COOH)CH_2CH_2N(CH_2CH_2SCH_2COOH)CH_2CH_2N(CH_2CH_2SCH_2COOH)CH_2-]$;

ethylenebis(carboxymethyliminomethylene) $[-CH_2N(CH_2COOH)CH_2CH_2N(CH_2COOH)CH_2-]$;

20 carboxymethyliminobis(methylene) $[-CH_2N(CH_2COOH)CH_2-]$;

or the salts of the exemplified carboxylic acid containing groups, including, for example, the metal and ammonium salts of such acids as described for R^2 above. In especially preferred embodiments, R^2 is

25 methyleneiminodiacetic acid or a salt thereof.

Each R^3 independently is selected from the groups specified for R. R^3 preferably represents hydrogen.

Each R^4 is independently selected from hydrogen or a protein reactive group.

In formula A-I above, n is 0, 1, 2, 3 or 4; m is 0 or 1; and o is 0 or 1; provided that at least one of n and m is 0.

At least one of the R, R^1 , R^3 and R^4 groups present is a protein reactive group. Preferably, no more than one of the R, R^1 , R^3 and R^4 groups on each aromatic ring is a protein reactive group. Most preferably, only one of the R, R^1 , R^3 and R^4 groups per molecule is a protein reactive group.

By "protein reactive group" it is meant any group which can react with any functional groups typically found on proteins. However, it is specifically contemplated that the protein reactive group can be conjugated to nonprotein biomolecules. Thus the protein reactive groups useful in the practice of this invention include those groups which can react with any biological molecule containing an immunoreactive group, whether or not the biological molecule is a protein to form a linking group between the complexing agent and the immunoreactive group.


Preferred protein reactive groups can be selected from but are not limited to: (1) A group that will react directly with the amine or sulfhydryl groups on the protein or biological molecule containing the immunoreactive group, for example, active halogen containing groups including, for example, chloromethylphenyl groups and chloroacetyl [$\text{Cl}-\text{CH}_2\text{CO}-$] groups, activated 2-leaving group substituted ethylsulfonyl and ethylcarbonyl groups such as 2-chloroethylsulfonyl and 2-chloroethylcarbonyl; vinylsulfonyl; vinylcarbonyl; epoxy; isocyanato; isothiocyanato; aldehyde; aziridine; succinimidoxycarbonyl; activated acyl groups such as carboxylic acid halides; mixed anhydrides and the like; and

other groups known to be useful in conventional photographic gelatin hardening agents.

- (2) A group that can react readily with modified proteins or biological molecules containing the immunoreactive group, i.e., proteins or biological molecules containing the immunoreactive group modified to contain reactive groups such as those mentioned in (1) above, for example, by oxidation of the protein to an aldehyde or a carboxylic acid, in which case the "protein reactive group" can be selected from amino, alkylamino, arylamino, hydrazino, alkylhydrazino, arylhydrazino, carbazido, semicarbazido, thiocarbazido, thiosemicarbazido, sulfhydryl, sulfhydrylalkyl, sulfhydrylaryl, hydroxy, carboxy, carboxyalkyl and carboxyaryl. The alkyl portions of the protein reactive group can contain from 1 to about 20 carbon atoms as described for R above. The aryl portions of the protein reactive group can contain from about 6 to about 20 carbon atoms as described for R above.
- (3) A group that can be linked to the protein or biological molecule containing the immunoreactive group, or to the modified protein as noted in (1) and (2) above by use of a crosslinking agent. Certain useful crosslinking agents, such as, for example, difunctional gelatin hardeners, bisepoxides and bisisocyanates become a part of, i.e., a linking group in, the protein-complexing agent conjugate during the crosslinking reaction. Other useful crosslinking agents, however, facilitate the crosslinking, for example, as consumable catalysts, and are not present in the final conjugate. Examples of such crosslinking agents are carbodiimide and carbamoylonyl crosslinking agents as disclosed in U.S. Patent 4,421,847, the disclosure of which is hereby incorporated by reference in its entirety, and the dication ethers of U.S. Patent 4,877,724, the disclosure of which is hereby

incorporated by reference in its entirety. With these crosslinking agents, one of the reactants must have a carboxyl group and the other an amine or sulfhydryl group. The crosslinking agent first reacts selectively with the carboxyl group, then is split out during reaction of the "activated" carboxyl group with an amine to form an amide linkage between the protein and metal complexing agents having the structure A-I above, this covalently bonding the two moieties. An advantage of this approach is that crosslinking of like molecules, e.g., proteins with proteins or complexing agents with complexing agents is avoided, whereas the reaction of difunctional crosslinking agents is nonselective and unwanted crosslinked molecules are obtained. Especially preferred protein reactive groups include amino and isothiocyanato.



Especially preferred complexing agents include species 1-58 set forth below.





* The hydrogen in all carboxy ($-\text{COOH}$) groups can be replaced by a metal or ammonium cation.

** All NH_2 groups can be replaced by NCS or $\text{NHCSN}(\text{CH}_3)\text{NH}_2$



2. H




the 2 R²s taken together =







3. H


the 2 R²s taken together =

4. H


the 2 R²s taken together =










the 2 R²s taken together =















the 2 R²s taken together =





the 2 R²s taken together =



















H the 2 R's taken together = H



B-4







the 2 R²s taken together =






SUBSTITUTE SHEET



37. -NCS

the 2 R²s taken together =the 2 R²s taken together =39. -NH₂the 2 R²s taken together =


40. -NCS

the 2 R²'s taken together =the 2 R²'s taken together =


B-5







 R^1





H

 R^2 






H











B-6





SUBSTITUTE SHEET





the 2 R²s taken together =





the 2 R²s taken together =





the 2 R²s taken together =




the 2 R²s taken together =



the 2 R²s taken together =




the 2 R²s taken together =



Preferred classes of complexing agents for use herein include terpyridines represented by structure A-III above and phenanthrolines represented by structure A-IV above. A particularly preferred class of complexing agents has the structure A-III above wherein $n=1$ and R is a substituted phenyl containing an alkyl or alkoxy substituent and a protein reactive group. Representative preferred species of complexing agents include compounds 20-32 depicted above. The currently most preferred complexing agent is TMT (compound 21).

This invention provides novel terpyridines having the structure A-III set forth in the summary above wherein $n=1$ and R is



R⁶ is a protein reactive group. R⁵ is alkyl, preferably containing from 1 to about 20, more preferably from 1 to 8 carbon atoms, such as methyl, ethyl and the like; or alkoxy, the alkyl portion of which contains from 1 to about 20, more preferably from 1 to 8 carbon atoms, such as methoxy, ethoxy and the like. R⁶ is a protein reactive group as described above. Preferred protein reactive groups include amino, alkylamino, arylamino, carbazido, semicarbazido, thiosemicarbazido, thiocarbazido, isocyanato and isothiocyanato. Especially preferred protein reactive groups include amino, isothiocyanato, and semicarbazido. Especially preferred species include TMT (compound 21) and compound 20 (THT) depicted above.


Preferred phenanthrolines according to this invention have the structure A-IV above wherein at least one R⁴ is a protein reactive group. Preferred



protein reactive groups include those specified for R⁶ above.

The polypyridine and phenanthroline complexing agents having metal complexing sites, e.g., heteroatoms and iminodiacetate groups can be prepared by techniques known in the art. Suitable reaction schemes are illustrated in U.S. Patent 4,837,169 and U.S. Patent 4,859,777, the disclosures of which are hereby incorporated by reference.



The preparation of currently preferred compounds of this invention, namely: 4'-(3-amino-4-methoxyphenyl)-6,6"-bis(N',N'-dicarboxymethyl-N-methylhydrazino)-2,2':6',2"-terpyridine, tetrasodium salt (THT) and 4'-(3-amino-4-methoxyphenyl)-6,6"-bis[N,N-di-(carboxymethyl)aminomethyl]-2,2':6',2"-terpyridine, tetrasodium salt (TMT) are illustrated in the following Reaction Scheme I:



Reaction Scheme I Synthesis of THT and TMT








THT Synthesis





TMT Synthesis





The addition to these molecules of the requisite protein reactive group described above can be accomplished by conventional chemical reactions. For example, amine groups can be added to polypyridines and phenanthrolines by nitration followed by reduction of the nitro groups to amines. If desired, the amine groups can be readily converted to isocyanate groups by reaction with phosgene to produce the carbamoyl chloride which, upon heating, releases HCl to produce the isocyanate. Carboxy groups can be added by treatment of the amine-substituted polypyridines and phenanthrolines with agents such as glutaric anhydride, followed by suitable selective activation of the carboxyl functionality. Cyclic acetal protected aldehydes can be carried through the reaction sequence necessary to synthesize the polypyridine chelators, and then deprotected before protein conjugation.

The class of terpyridines conforming to Structure A-III above and containing a phenyl group substituted with an alkyl or alkoxy substituent and a protein reactive group is particularly advantageous from a synthetic standpoint. The presence of the alkyl or alkoxy group on the dibrominated starting material

(3) provides enhanced solubility in THF which is a preferred solvent used in the preparation of the intermediate diol (6).

5 The novel terpyridines and phenanthrolines of this invention can be prepared according to the above described synthetic techniques. Additional illustrative preparations are set forth in the examples which follow.

10 The targeting radioactive immunoreagent of this invention includes a radionuclide ion. The radionuclide ion can be selected, for example, from Sc, Fe, Pb, Ga, Y, Bi, Mn, Cu, Cr, Zn, Ge, Mo, Tc, Ru, In, Sn, Sm, Sb, W, Re, Po, Ta and Tl ions. Preferred radionuclides include $^{44}\text{Sc}^{+++}$, $^{64,67}\text{Cu}^{++}$, $^{111}\text{In}^{+++}$,
15 $^{212}\text{Pb}^{++}$, $^{68}\text{Ga}^{++}$, $^{90}\text{Y}^{+++}$ and $^{212}\text{Bi}^{+++}$ ions. Of these, the most preferred are $^{90}\text{Y}^{+++}$ ions.

The metal radionuclide ion and the complexing agent are easily complexed by merely mixing an aqueous solution of the complexing agent with a metal
20 radionuclide salt in an aqueous solution preferably having a pH of 4 to 11. The salt can be any water soluble salt of the metal such as halogen salts. The chelate is generally prepared in aqueous solution at a pH of between 5 and 9 and preferably 6 to 8. The
25 complex optionally is mixed with buffers such as acetate, phosphate and borate to produce the optimum pH.

The targeting immunoreagent of this invention includes an immunoreactive group covalently bonded to
30 the complexing agent. The targeting immunoreagent thus comprises a conjugate of a complex having the structure A-I above and the immunoreactive group. The complexing agent and the metal radionuclide can be complexed either before or after the complexing agent is attached
35 to the immunoreactive group. As used herein the term "immunoreactive group" is meant to include any organic

compound which is capable of covalently bonding to the complexing agent and which is found in a living organism or is useful in the diagnosis, treatment or genetic engineering of cellular material or living organisms, and which has a capacity for interaction with another component which may be found in biological fluids or associated with cells to be treated such as tumor cells.

Depending upon the intended use, the immunoreactive group can be selected from a wide variety of naturally occurring or synthetically prepared materials, including, but not limited to enzymes, amino acids, peptides, polypeptides, proteins, lipoproteins, glycoproteins, hormones, drugs (for example digoxin, phenytoin, phenobarbitol, thyroazine, triiodothyronine, gentamicin, carbamazepine, and theophylline), steroids, vitamins, polysaccharides, viruses, protozoa, fungi, parasites, rickettsia, molds, and components thereof, blood components, tissue and organ components, pharmaceuticals, haptens, lectins, toxins, nucleic acids (including oligonucleotides), antibodies, antigenic materials (including proteins and carbohydrates), avidin and derivatives thereof, biotin and derivatives thereof, and others known to one skilled in the art.

Preferred immunoreactive groups for use in the practice of this invention are those which have a receptor molecule specific to a ligand of interest. Thus, a specific binding reaction involving the reagent can be used for the targeting expected. Examples of such ligand-receptor complexes include, but are not limited to antibody-antigen, avidin-biotin, repressor (inducer) - promoter of operons and sugar-lectin complexes. Additionally, complementary nucleic acids, i.e., a hybridized product of complementary strands, are also considered specific binding materials as the

term is used herein.

Useful immunoreactive groups include (1) any substance which, when presented to an immunocompetent host, will result in the production of a specific antibody capable of binding with that substance, or (2) the antibody so produced, which participates in an antigen-antibody reaction. Thus, the immunoreactive group can be an antigenic material, an antibody, or an anti-antibody. Both monoclonal and polyclonal antibodies are useful. The antibodies can be whole molecules or various fragments thereof, as long as they contain at least one reactive site for reaction with the reactive groups on the complexing agent or with linking groups as described herein.

In certain embodiments, the immunoreactive group can be an enzyme which has a reactive group for attachment to the complexing agent. Representative enzymes include, but are not limited to, aspartate aminotransaminase, alanine aminotransaminase, lactate dehydrogenase, creatine phosphokinase, gamma glutamyl transferase, alkaline acid phosphatase, prostatic acid phosphatase, horseradish peroxidase and various esterases.

If desired, the immunoreactive group can be modified or chemically altered to provide reactive groups for attaching to the complexing agent by techniques known to those skilled in the art. Such techniques include the use of linking moieties and chemical modification such as described in WO-A-89/02931 and WO-A-89/2932, which are directed to modification of oligonucleotides, and U.S. Patent 4,719,182 the disclosures of which are hereby incorporated by reference in their entirety.

Two highly preferred uses of the targeting immunoreagents of this invention are for the diagnostic imaging of tumors and the radiological treatment of

tumors. Preferred immunological groups therefore include antibodies to tumor-associated antigens. Specific examples include B72.3 antibodies (described in U.S. Patent Nos. 4,522,918 and 4,612,282) which
5 recognize colorectal tumors, 9.2.27 anti-melanoma antibodies, D612 antibodies which recognize colorectal tumors, UJ13A antibodies which recognize small cell lung carcinomas, NRLU-10 antibodies which recognize small cell lung carcinomas and colorectal tumors (Pan-
10 carcinoma), 7E11C5 antibodies which recognize prostate tumors, CC49 antibodies which recognize colorectal tumors, TNT antibodies which recognize necrotic tissue, PR1A3 antibodies, which recognize colon carcinoma, ING-1 antibodies, which are described in International
15 Patent Publication WO-A-90/02569, B174 antibodies which recognize squamous cell carcinomas, B43 antibodies which are reactive with certain lymphomas and leukemias and others which may be of particular interest.

Such antibodies and other useful immunological
20 groups described above are large, complex molecules having multiple sites for appendage of the complexing agent. Consequently, the immunoreactive group can have appended to it additional complexing agents via one of the protein reactive groups. Thus, the term
25 immunoreactive group is intended to include immunological groups having complexing agent molecules bonded thereto through one or more protein reactive groups.

Additionally, an antibody or fragment thereof
30 containing a carbohydrate region can be attached to the complexing agent through the carbohydrate region of the antibody, such as described in U.S. Patent 4,937,183, the disclosure of which is hereby incorporated by reference in its entirety. Useful methods for
35 attaching an antibody are also described in U.S. Patent Nos. 4,671,958; 4,699,784; 4,741,900; and 4,867,973.

The term "protein reactive group" as defined herein is intended to include such linkages.

Other techniques for performing the covalent binding of the immunoreactive group to the radioactive metal complexing agents are known in the art and include simply mixing the materials together.

The radioactive immunoreagent of this invention can contain any ratio of metal radionuclide ion to complexing agent. In preferred embodiments, the mole ratio of metal ion to complexing agent is from about 1:100 to about 1:1.

The ratio of chelate to immunoreactant can vary widely from about 0.5:1 to 10:1 or more. In some embodiments, the mole ratio of chelate to immunoreactive groups is from about 1:1 to about 6:1.

Figs. 6 and 7 are spectra of complexing agents and metal complexes of this invention. Portions of the spectra do not overlap with those of the proteins to which the chelating agents are chemically bonded. Similar spectral shifts were obtained between chelating agents of this invention and other representative cations such as Ga^{+3} , Bi^{+3} , In^{+3} , Sc^{+3} and Cu^{+2} . Thus, the immunoreagent of this invention can be readily spectrophotometrically analyzed.

The following examples further illustrate the invention:

Preparation 1 Preparation of 4'-(3-Amino-4-methoxy-phenyl)-6,6"-(N,N-di(carboxymethyl)amino-methyl)-2,2':6',2"-terpyridine.
Tetrasodium Salt (TMT)

Part A - Pyridinium Bromide 1

2-Acetyl-6-bromopyridine was synthesized by the method of J.E. Parks, B.E. Wagner, and R.E. Holm, J. Organometal. Chem 56, 53-66 (1973). 2-Acetyl-6-bromopyridine (20.0 g, 100 mmol) was treated with bromine (6.2 mL, 0.12 mol) at reflux in 200 mL of $CHCl_3$.

for 45 min. The solution was cooled to room temperature then washed with dilute aqueous $\text{NaHCO}_3/\text{Na}_2\text{S}_2\text{O}_3$. The organic phase was dried over Na_2SO_4 , filtered, and evaporated to give an oil. The oil was dissolved in 200 mL of THF and 30 mL of pyridine was added. The resulting solution was refluxed for 30 min. The mixture was cooled and filtered to give 26.1 g of off-white powder (73%): mp 256°C dec (discolors at 245°C). Anal. Calcd. for $\text{C}_{12}\text{H}_{10}\text{Br}_2\text{N}_2\text{O}$: C, 40.26; H, 2.82; N, 7.82. Found: C, 40.12; H, 2.85; N, 7.79. The NMR and IR spectra were consistent with the assigned structure and the product was homogeneous by TLC.

Part B - Chalcone 2

Potassium hydroxide (18.2 g, 325 mmol) was dissolved in 100 mL of H_2O , and 100 mL of methanol was added. 2-Acetyl-6-bromopyridine (65 g, 325 mmol) and (68 g, 650 mmol) of p-anisaldehyde were dissolved together in 400 mL of methanol, and the solution was poured into the KOH solution. Precipitation of product began within a few minutes, and the reaction was allowed to stand at room temperature overnight. The precipitate was collected, washed with isopropanol, and dried to yield 79 g (76%) of the product as a yellow solid, mp $100\text{--}102^\circ\text{C}$. FDMS (m/e) 317 M. An aliquot was purified by column chromatography on Woelm Silica gel, elution with 100% dichloromethane. Anal. Calcd for $\text{C}_{15}\text{H}_{12}\text{BrNO}_2$: C, 56.63; H, 3.80; N, 4.40. Found: C, 56.66; H, 3.87; N, 4.41. The NMR and IR spectra were consistent with the assigned structure and the product was homogeneous by TLC.

Part C - Dibromoterpyridine 3

Pyridinium bromide 1 (11.3 g, 31.6 mmol) and

chalcone 2 (10.0 g, 31.4 mmol) were refluxed in 100 mL of AcOH with 10 g of NH_4OAc for 16 hours. The

solution was cooled and filtered, and the solid was washed with AcOH then EtOH to give 13.48 g of white

5 crystals (86%): mp 203–204.5°C. FDMS (m/e) 495 (M+). Anal. Calcd for $\text{C}_{22}\text{H}_{15}\text{Br}_2\text{N}_3\text{O}$: C, 53.1; H, 3.0; N, 8.5.

Found: C, 52.9; H, 3.1; N, 8.4. The NMR and IR spectra were consistent with the assigned structure and the product was homogeneous by TLC.

10 Part D - Terpyridinediol 6

Dibromide 3 (7.46 g, 15.5 mmol) in 100 mL of dry THF was added dropwise to a solution of 28.1 mL of 1.6 M n-BuLi in 20 mL of dry THF under N_2 over a 12 min period. The temperature was maintained below -75°C

15 during the addition with a dry ice/acetone bath. The resulting dark green solution was stirred for 10 minutes followed by addition of 7.5 mL of dry DMF over a 2 min period. After 10 min, 90 mL of a 10% HCl solution was added and the resulting solution was

20 stirred for 45 min with continued cooling. The mixture was partitioned between CHCl_3 and H_2O (both solvents pre-cooled to 4°C) in a separatory funnel. The phases were shaken frequently and allowed to stand at ambient temperature for 15–30 minutes, until the color of the
25 organic phase gradually changed from a greenish hue to golden yellow. The organic phase was washed with sat. NaCl then evaporated to leave a cream-colored residue.

This material was triturated with CH_3CN to yield the product as an off-white solid (3.53 g, 60%), mp 225-227°C. FDMS (m/e) 395M. Anal. Calcd for $\text{C}_{24}\text{H}_{17}\text{N}_3\text{O}_3$: C, 72.90; H, 4.33; N 10.63. Found: C, 72.44; H, 4.31; N, 10.46.

The crude dialdehyde (3.53 g, 8.93 mmol) was refluxed with 1 g of NaBH_4 in a mixture of 70 mL of THF and 70 mL of abs. EtOH for 15 min under N_2 . After concentration in vacuo, the residue was refluxed for 30 min in dilute NaHCO_3 , cooled, filtered, washed with H_2O , then dried to give diol 6 as a white solid (3.35 g, 94.4%). mp 187-189°C. FDMS (m/e) 400 MH^+ , 399 M. Anal. Calcd for $\text{C}_{24}\text{H}_{21}\text{N}_3\text{O}_3$: C, 72.17; H, 5.30; N, 10.52. Found: C, 71.71; H, 5.20; N, 10.37. The NMR and IR spectra were consistent with the assigned structure and the product was homogeneous by TLC.

Part E - Tetraester 7

Diol 6 (15.4 g, 38.5 mmol) was suspended in a mixture of 17 mL of Et_3N in 175 mL of CH_2Cl_2 with stirring at 8°C. To this suspension, a solution of $(\text{CH}_3\text{SO}_2)_2\text{O}$ (16.8 g, 96.5 mmol) in 50 mL of CH_2Cl_2 was added dropwise over a 10 min period. The reaction mixture was shaken with water. The organic layer was dried over Mg_2SO_4 , filtered, and concentrated nearly to dryness. Addition of EtOAc produced the bismesylate as white crystals which were collected and dried (17.2 g, 80.4%). A mixture of the bismesylate (0.50 g, 0.96

mmol), diisopropylethylamine (0.26 g, 2.0 mmol), and diethyliminodiacetate (0.38 g, 2.0 mmol) was stirred for 16 hours in 20 mL of dry DMF. The mixture was concentrated in vacuo and the residue was partitioned between Et₂O and H₂O. The Et₂O phase was washed two additional times with water then dried over Na₂SO₄ and evaporated to give the product as a pale yellow oil (0.58 g, 82%). Anal. calcd for C₄₀H₄₇N₅O₉: C, 64.76; H, 6.39; N, 9.44. Found: C, 64.35; H, 6.17; N, 9.39. The NMR and IR spectra were consistent with the assigned structure and the product was homogeneous by TLC.

Part F - Nitrotetraester 8

Terpyridine tetraester 7 (3.27 g, 4.41 mmol) was dissolved in 60 mL of conc. H₂SO₄ to give a red-orange solution. The mixture was cooled to 0°C and a 1/10 (v/v) mixture of conc. HNO₃ in conc. H₂SO₄ was added such that 4.41 mmol of HNO₃ was delivered. The color of the solution turned pale yellow after addition was completed. The reaction mixture was stirred for 15 min at 0°C, then poured onto crushed ice. Dilute K₂CO₃ was added until pH 8 was reached. The aqueous solution was extracted three times with CH₂Cl₂. The organic layers were combined, dried over Na₂SO₄, and concentrated to give a yellow oil, which was chromatographed on silica gel (5% MeOH/CH₂Cl₂). The fractions containing product were combined and

evaporated to give the product which was recrystallized three times from MeOH to yield an off-white solid (1.84 g, 53%): mp 76-79°C. FDMS (m/e) 787 MH^+ , 786 M. Anal. calcd for $C_{40}H_{46}N_6O_{11}$: C, 61.06; H, 5.89; N, 10.68. Found: C, 60.69; H, 6.22; N, 11.04. The NMR and IR spectra were consistent with the assigned structure and the product was homogeneous by TLC.

Part G - Aminoterpyridine Tetraester 9

10 Nitrotetraester 8 (1.80 g, 2.29 mmol) was dissolved in a mixture of 90 mL of THF and 90 mL of abs. EtOH. Ammonium formate (2.89 g, 45.8 mmol) dissolved in 16 mL of H_2O was added, followed by 4.8 g of 10% Pd/C (4.6 mmol). After stirring at room
15 temperature for 2 h, the reaction was filtered through a diatomaceous earth filter pad. The filter pad was washed well with THF, abs. EtOH, and CH_2Cl_2 . The filtrate was concentrated, and the residue was partitioned between CH_2Cl_2 and aqueous NaCl. The
20 organic phase was concentrated then purified on SiO_2 with 10% MeOH/ $CHCl_3$ to give the product as a straw-colored oil (0.80 g, 46%). FDMS (m/e) 757 MH^+ , 756 M. Anal. calcd for $C_{40}H_{48}N_6O_9 \cdot 1/2 H_2O$: C, 62.73; H, 6.45; N, 10.97. Found: C, 62.98; H, 6.47; N, 10.67. The
25 NMR and IR spectra were consistent with the assigned structure and the product was homogeneous by TLC.

Part H - Aminotetraacid 10 (TMT)

Amine tetraester 9 (0.75 g, 0.99 mmol) was

stirred with 4 equiv of NaOH in a mixture of 50 mL of MeOH and 2 mL of H₂O for 16 hours at room temperature.

The mixture was concentrated to give the product as a solid dihydrate (0.72 g, 94%): FABMS m/e 640 (M⁺ for tetracarboxylate). Anal. calcd. for C₃₂H₂₈N₆Na₄O₉·2H₂O: C, 50.01; H, 4.20; N, 10.93.

Found: C, 49.82; H, 4.12; N, 10.74. The NMR and IR spectra were consistent with the assigned structure and the product was homogeneous by TLC.

10 Preparation 2 Preparation of 4'-(3-amino-4-methoxy-phenyl)-6,6"-bis(N'N'-dicarboxymethyl-N-methylhydrazino)-2,2':6',2"-terpyridine, tetrasodium salt (THT)

15 Part A - 6,6"-Bis(N-methylhydrazino)-4'-(4-methoxy-phenyl)-2,2':6',2"-terpyridine
6,6"-Dibromo-4'-(4-methoxyphenyl)-

2,2':6',2"-terpyridine (1.0 g, 2 mmol) was refluxed in 20 mL of methylhydrazine for 16 hours under nitrogen. The solution was cooled, and the resulting precipitate filtered, and dried to a constant weight to give 0.68 g of cream-colored solid, mp 218-220°C. FDMS (m/e) 428 MH⁺, 427M⁺. Anal. calcd for C₂₄H₂₅N₇O·0.25 H₂O:

20 C, 66.72; H, 5.96; N, 22.70. Found: C, 66.85; H, 5.85; N, 23.0. The NMR and IR spectra were consistent with the assigned structure.

25 Part B - 6,6"-Bis(N',N'-di(ethoxycarbonylmethyl)-N-methylhydrazino)-4'-(4-methoxyphenyl)-2,2':6',2"-terpyridine

The bis(methylhydrazine)terpyridine of Part A (3.50 g, 82 mmol), ethyl bromoacetate (13.2 mL, 820 mmol), 2,6-lutidine (9.6 mL, 820 mmol), and sodium iodide (0.35 g, 2 mmol) were added to 350 mL of acetonitrile, and the solution was refluxed under N₂

for 48 hours, when an additional 4.1 mL (370 mmol) of ethyl bromoacetate and 4.8 mL (410 mmol) of 2,6-lutidine were added. The reaction solution was refluxed for an additional 48 hours, and cooled. A
5 copious amount of white salt resulting from excess bromoacetate and lutidine was filtered and discarded. The filtrate was concentrated, and the concentrated material was dissolved in dichloromethane, and extracted two times with dilute aqueous sodium
10 chloride. The organic phase was concentrated under high vacuum until free of odors of bromoacetate and lutidine. The crude oil was purified on a Woelm silica gel column (36 x 2 in.). The column was eluted initially with 100% dichloromethane followed by 50/1
15 dichloromethane/acetone, with gradual increase of the concentration of acetone to 25/1 dichloromethane/acetone. Concentration of purified fractions gave 2.91 g (46%) of light straw-colored oil. A fraction of the purified oil, upon standing at room
20 temperature, crystallized, and after trituration with methanol gave a white solid, mp 100-103°C. Anal. calcd for $C_{40}H_{49}N_7O_9$: C, 62.24; H, 6.40; N, 12.70. Found: C, 62.31; H, 6.32; N, 12.69. The NMR and IR spectra were consistent with the assigned structure, and the
25 product was homogeneous by TLC.

Part C - 6,6"-Bis(N',N'-di(ethoxycarbonylmethyl)-N-methylhydrazino)-4'-(4-methoxy-3-nitrophenyl)-

2,2':6',2"-terpyridine.

The terpyridine tetraester of Part B (0.659 g, 0.84 mmol) was dissolved in 5 mL of conc H_2SO_4 to give a red-orange solution. The mixture was cooled to 0°C , and a mixture of conc HNO_3 in conc H_2SO_4 was added such that 0.84 mmol of HNO_3 was delivered. The color of the solution turned pale yellow after addition was completed. The reaction mixture was stirred for 15 minutes at 0°C , then poured onto crushed ice. Dilute K_2CO_3 was added until pH 8 was reached. The aqueous solution was extracted three times with CH_2Cl_2 . The organic layers were combined and dried over Na_2SO_4 , and concentrated to give a yellow oil, which was chromatographed on silica gel (20/1 methylene chloride/acetone). The fractions containing product were combined and evaporated to give the product, a pale yellow glass. Yield 0.15 g (22%). Anal. calcd for $\text{C}_{40}\text{H}_{48}\text{N}_8\text{O}_{11}$: C, 58.82; H, 5.92; N, 13.72. Found: C, 58.76; H, 5.61; N, 13.84. FDMS (m/e) 816 M. The NMR and IR spectra were consistent with the assigned structure, and the product was homogeneous by TLC.

Part D - 4'-(3-Amino-4-methoxyphenyl)-6,6"-bis(N',N'-di(ethoxycarbonylmethyl)-N-methylhydrazino)-2,2':6',2"-terpyridine

The nitrotetraester of Part C (0.72 g, 0.88 mmol) was dissolved in a mixture of 30 mL of THF and 30 mL of ethanol. Ammonium formate (1.08 g, 17.1 mmol)

dissolved in 6 mL H_2O was added, followed by 1.8 g of
10% Pd/C (1.7 mmol). After stirring at room
temperature overnight, the reaction was filtered
through a diatomaceous earth filter pad. The filter
5 pad was washed well with THF, absolute EtOH, and
 CH_2Cl_2 . The filtrate was concentrated, and the residue
was dissolved in chloroform, and the chloroform
solution was washed two times with water. The organic
phase was concentrated to an oil. Addition of methanol
10 to the oil resulted in crystallization of the oil. The
material was slurried in 20 mL of methanol, and
filtered, and air dried to yield 0.64 g of cream
colored solid (92.7%). Anal. calcd for $C_{40}H_{50}N_8O_9$: C,
61.06, H, 6.40; N, 14.24. Found: C, 60.74; H, 6.39;
15 N, 14.01. The NMR and IR spectra were consistent with
the assigned structure and the product was homogeneous
by TLC.

Part E - 4'-(3-Amino-4-methoxyphenyl)-6-6"-bis(N',N'-
dicarboxymethyl-N-methylhydrazino)-2,2':
20 6',2"-terpyridine, tetrasodium salt (THT)

The amine tetra ester of Part D (0.60 g, 0.76
mmol) was stirred with 4 equivalents of NaOH in a
mixture of 25 mL of MeOH and 1 mL of H_2O for about 16
hrs at room temperature. The mixture was concentrated
25 to give a quantitative yield of solid tetracarboxylate.
Anal. calcd for $C_{32}H_{30}N_8Na_4O_9 \cdot 2H_2O$: C, 48.12; H, 4.29;
N, 14.03. Found: C, 48.01; H, 3.97; N, 12.77.

Preparation 3 Preparation of 6,6''-Bis(N',N'-di-carboxymethyl-N-methylhydrazino)-4'-(3-isocyanato-4-methoxyphenyl)-2,2':6',2''-terpyridine, Sodium Salt

5 The THT amine tetrasodium salt of Preparation 2, Part E (0.39 g, 0.51 mmol) was dissolved in 80 mL of methanol. At room temperature, 0.58 g (0.50 mmol) of thiophosgene in 1.0 mL tetrahydrofuran was added, followed by addition of 0.51 g (0.50 mmol) of
10 triethylamine in 1.0 mL of tetrahydrofuran. The reaction was then concentrated to a residue, which was slurried in dichloromethane, and filtered to yield 0.26 g (65%) of yellow solid. The infrared spectrum was consistent with the assigned structure. The product
15 was believed to be the monocarboxylic acid trisodium salt.

Preparation 4 Preparation of 5-Amino-2,9-Bis[N,N-di-(carboxymethyl)aminomethyl]-1,10-phenanthroline, Tetrasodium Salt

20 Part A - 2,9-Dimethyl-5-nitro-1,10-phenanthroline
 Neocuproine hydrochloride, hemihydrate (25.0 g, 99 mmol) was dissolved in 100 mL of concentrated nitric acid. 200 mL of concentrated sulfuric acid was added, and the reaction was heated at reflux for 2.5
25 hours, and then allowed to stand at room temperature for 8 days. The reaction mixture was then gradually added to a mixture of about 3 Kg of ice and 351 g (8.8 mole) of LiOH·H₂O, while stirring with a glass rod.
During the neutralization procedure, ice was added as
30 necessary so that unmelted ice was always present during the neutralization, and the neutralization reaction was also simultaneously cooled by an

acetone/ice bath. After the addition was completed, the pH of the reaction mixture was 12. The aqueous mixture was extracted two times with methylene chloride, first with 1000 mL, and then with 400 mL.

5 The methylene chloride fractions were combined, and extracted once with 500 mL of distilled H₂O. The methylene chloride phase was concentrated to a residue, and triturated with 500 mL of acetonitrile. The solid was filtered, washed with acetonitrile, and dried under
10 vacuum to give 10.3 g of cream yellow solid, which was dissolved in 100 mL of refluxing acetonitrile, allowed to crystallize at room temperature, and was filtered to give 8.5 g (34%) of cream-yellow crystals, mp 184-186°C. FDMS (m/e) 253M. Anal. calcd for
15 C₁₄H₁₁N₃O₂·0.25 H₂O: C, 65.23; H, 4.50; N, 16.30. Found: C, 65.57; H, 4.45; N, 16.27. The NMR and IR spectra were consistent with the assigned structure and the product was homogeneous by TLC.

Part B - 2,9-Di(bromomethyl)-5-nitro-1,10-phenanthroline
20

2,9-Dimethyl-5-nitro-1,10-phenanthroline (1.0 g, 4 mmol) and N-Bromosuccinimide (1.42 g, 8 mmol) were mixed together into a homogeneous mixture in a beaker, and added all at once to 20 mL of refluxing (181°C) o-dichlorobenzene. The heating was continued at the
25 reflux temperature for 2 minutes, and then the reaction was allowed to cool to room temperature. The reaction

mixture was purified on a column (18 x 2 in.) of Woelm silica gel, by elution with 100% dichloromethane, to give 0.47 g (29%) of purified material, a yellow oil. Rf on TLC, 0.5 (40/1 CH₂Cl₂/acetone). Trituration of

5 an aliquot of the purified product gave a gray-white solid, which decomposed at 146°C. Anal. calcd for C₁₄H₉Br₂N₃O₂•0.5 H₂O: C, 40.02; H, 2.39; N, 10.00.

Found: C, 40.36; H, 2.67; N, 9.77. FDMS (m/e) 409 M.

The NMR and IR spectra were consistent with the

10 assigned structure.

Part C - 2,9-Bis[N,N-di(ethoxycarbonylmethyl)-aminomethyl]-5-nitro-1,10-phenanthroline

2,9-Di(bromomethyl)-5-nitro-1,10-

15 phenanthroline (2.28 g, 5.5 mmol) and 2.08 g (11 mmol) of 1,8-bis(dimethylamino)naphthalene were dissolved together in 25 mL of 1-methyl-2-pyrrolidinone, and 2.08 g (11 mmol) of diethyl iminodiacetate was added. The reaction was sealed with a ground glass stopper, and

20 after approximately 10 minutes of stirring at room temperature, white precipitate began forming. The reaction was then placed in the refrigerator (at 4°C) and stored for about 20 hours. The reaction mixture was partitioned between 400 mL of diethyl ether and 400

25 mL of distilled H₂O and the ether fraction was

extracted a total of 5 times with 400 mL portions of distilled H₂O. The ether phase was concentrated and

dissolved in 400 mL of methylene chloride. The methylene chloride phase was extracted once with distilled H₂O, dried with a mixture of Celite diatomaceous earth and sodium sulfate, and then filtered, and concentrated to a dark oil. A column of Woelm silica gel was prepared in 10/1 methylene chloride/methanol (20 inches in height by 2 inches in diameter), and the crude oil was applied to the column in minimal methylene chloride. About 50 mL of methylene chloride was applied to the column, and then the column was eluted with 10/1 CH₂Cl₂/methanol. Fractions (100 mL each) were collected, and the fractions of pure product by TLC (10/1 methylene chloride/methanol, R_f 0.4) were combined and concentrated to yield 1.17 g (34%) of product, an oil of reddish-amber hue. FDMS (m/e) 628 MH⁺. Anal. calcd for C₃₀H₃₇N₅O₁₀·2 H₂O: C, 54.29; H, 6.23; N, 10.55. Found: C, 54.71; H, 5.58; N, 10.53. The NMR and IR spectra were consistent with the assigned structure.

Part D - 5-Amino-2,9-bis[N,N-di(ethoxycarbonylmethyl)-aminomethyl]-1,10-phenanthroline

2,9-Bis[N,N-di(ethoxycarbonylmethyl)amino-methyl]-5-nitro-1,10-phenanthroline (3.37 g, 5.4 mmol) was dissolved in a solution of 100 mL tetrahydrofuran and 200 mL absolute ethanol. A solution of 6.8 g (108 mmol) of ammonium formate in 22 mL of distilled H₂O was added, followed by 5.72 g of Palladium on carbon (10%)

(50% wet with water for safety). The reaction was stoppered with a gas bubbler, stirred at room temperature for 30 minutes, and then filtered through a diatomaceous earth filter pad. The solvents were then removed by a rotary evaporator. The residue was dissolved in 300 mL of dichloromethane, which was then extracted with 300 mL of distilled water. The organic phase was then extracted with 300 mL of saturated sodium chloride, then dried with a mixture of Celite diatomaceous earth and sodium sulfate, filtered, and concentrated under high vacuum to a dark amber oil which weighed 2.90 g. The oil was treated with 9.0 mL of propionitrile, which resulted in the crystallization of a yellow solid. The solid was filtered, and washed with 13 mL of propionitrile, and finally washed with 15 mL of hexanes, and air dried to a constant weight of 1.04 g (32%) mp 137-139°C. FDMS (m/e) 598 MH⁺, 597 M. Anal. calcd for C₃₀H₃₉N₅O₉·H₂O: C, 58.52; H, 6.71; N, 11.38. Found: C, 58.28; H, 6.55; N, 11.53. The NMR and IR spectra were consistent with the assigned structure.

Part E - 5-Amino-2,9-bis[N,N-di(carboxymethyl)aminomethyl]-1,10-phenanthroline, Tetrasodium Salt
5-Amino-2,9-bis[N,N-di(ethoxycarbonylmethyl)aminomethyl]-1,10-phenanthroline (0.90 g, 1.5 mmol) was dissolved in 200 mL of methanol. Sodium hydroxide (0.25 g, 6.25 mmol) dissolved in 10.0 mL of distilled H₂O was

added. The reaction was stoppered and stirred by magnetic stirring for 24 hours at room temperature. The solvent was removed by rotary evaporation, and the solid residue was triturated with methylene chloride and filtered to yield 0.81 g (94%). The tetrasodium salt was hygroscopic, and a significant amount remained adhered to the glass wall of the filtration funnel. Anal. calcd for $C_{22}H_{19}Na_4N_5O_8 \cdot 3 H_2O$: C, 42.11; H, 4.02; N, 11.16. Found: C, 42.01; H, 3.71; N, 11.03. The IR spectrum and FAB mass spectrum were consistent with the assigned structure.

Preparation 5 Preparation of 2,9-Bis[N,N-di(carboxymethyl)aminomethyl]-5-isothiocyanato-1,10-phenanthroline, Sodium Salt

5-Amino-2,9-bis[N,N-di(carboxymethyl)-aminomethyl]-1,10-phenanthroline, tetrasodium salt (0.16 g, 0.28 mmol) was dissolved in a solvent mixture of 20 mL methanol/4 mL distilled H_2O . Thiophosgene (0.28 mmol, i.e., 1.0 mL of a solution of 0.32 g thiophosgene in 10.0 mL THF) was added, followed immediately by addition of 0.28 mmol of triethylamine (1.0 mL of a solution of 0.28 g of triethylamine in 10.0 mL THF). The reaction was carried out at room temperature, and was then immediately concentrated to a residue, 0.16 g (94%) of yellow solid. The IR spectrum was consistent with the assigned structure. The product was believed to be the monocarboxylic acid

trisodium salt.


Preparation 6 Preparation of 6,6"-[N,N-di(carboxy-
methyl)aminomethyl]-4'-(3-isothio-
cyanato-4-methoxyphenyl)-2,2'-6',2"-
terpyridine, Tetrasodium Salt

5 The amino sodium tetracarboxylate of
 Preparation 1 (0.31 g, 0.42 mmol) was dissolved in 30
 mL of methanol, 1.00 mL of THF containing 0.42 mmol of
 thiophosgene was added and then 1.00 mL of THF
 10 containing 0.42 mmol of triethylamine was also added.
 The reaction solution was immediately concentrated to a
 residue triturated with triethylamine, the solid
 collected by filtration and washed with dichloromethane
 to give 0.30 g of product. The IR spectrum was
 15 consistent with the assigned structure.


Preparation 7. Trisodium 15-amino-3,5,6,8,9,11-hexahydro-
4,7,10-tris(carboxymethyl)-2,17:12,14-dietheno-1,4,7,10,13-
pentaazabenzocyclopentadecine

20

A




Part A. 3,5,6,8,9,11-Hexahydro-15-nitro-4,7,10-tris(p-
 toluenesulfonamido)-2,17:12,14-dietheno-1,4,7,10,13-
 25 pentaazabenzocyclopentadecine



100 mL of N-methylpyrrolidin-2-one containing 2 % by weight of 2,9-bisbromomethyl-5-nitro-1,10-phenanthroline prepared in Preparation 4, Part B is added via a syringe pump over 2 hours into 20 mL of magnetically stirred N-methylpyrrolidin-2-one which is held under argon at 20°C. Simultaneously, a 2 % solution in N-methylpyrrolidin-2-one of one equivalent of the disodium salt of diethylenetriamine N,N',N''-tri-p-toluenesulfonamide is added. The reaction is stirred for 6 additional hours after the addition at 20°C, 200 mL of the solvent is then distilled under high vacuum, the residual solution is cooled and added to 100 ml of ice water. The resulting precipitate is isolated by filtration, washed with cold water, and triturated with acetonitrile to provide the crude tri-p-toluenesulfonamide derivative.

Part B. 3,4,5,6,7,8,9,10,11-Nonahydro-15-nitro-2,17:12,14-dietheno-1,4,7,10,13-pentaazabenzocyclopentadecine




20

One part of crude macrocycle prepared in Part A is dissolved in 10 parts of concentrated sulfuric acid (96 %), and the reaction mixture is stirred and heated to 110°C under argon. After 24 hours, the solution is cooled with vigorous stirring in an ice water bath and is treated with 10 % sodium hydroxide solution until the pH reaches 10. The aqueous phase is extracted five times with equal volumes of chloroform, the extracts are combined and are dried over anhydrous sodium sulfate. The salts are then removed by filtration, and the filtrate is evaporated under

argon. The crude triamine can be purified by crystallization from ethanol/water.


5 Part C. 3,5,6,8,9,11-Hexahydro-15-nitro-4,7,10-tris(ethoxycarbonylmethyl)-2,17:12,14-dietheno-1,4,7,10,13-pentaazabenzocyclopentadecine



followed by 2 equivalents of 10% Pd/C. The reaction mixture is stirred at room temperature overnight under argon, the reaction mixture is filtered under argon, and the filtered catalyst is washed well with THF, ethanol, and then dichloromethane under argon. The combined filtrate and washings are concentrated, the residue is dissolved in chloroform, and the chloroform solution is washed twice with water. The organic phase is concentrated to an oil, and crystallization is promoted by the addition of methanol. The product is triturated in methanol, filtered, and then dried in air.

Part E. The amine triester of Part D is stirred at room temperature with three equivalents of sodium hydroxide in 5% aqueous methanol for 24 h. The solvent is removed under vacuum and the product (A) is triturated with a little cold methanol. The product A can be isolated by filtration.

Preparation 8. Trisodium 15-isothiocyanato-4,7,10-tris(carboxymethyl)-2,17:12,14-dietheno-1,4,7,10,13-pentaazabenzocyclopentadecine



25

One part by weight of the amine trisodium salt from Preparation 7, Part E, is stirred in 200 parts of methanol at room temperature.

This is treated at ambient temperature with one equivalent of thiophosgene dissolved in two parts by weight of tetrahydrofuran followed by one equivalent of triethylamine


30

dissolved in two parts by weight of tetrahydrofuran. After one hour, the reaction mixture is concentrated to a residue, slurried in dichloromethane, and the product is isolated by filtration.

5

Preparation 9. Trisodium 2-Methyl-3-thio-4-{15-[3,5,6,8,9,11-hexahydro-4,7,10-tris(carboxymethyl)-2,17:12,14-dietheno-1,4,7,10,13-pentaazabenzocyclopentadecynyl]}-semicarbazide

10




15

20

To a freshly prepared and stirred mixture of one part by weight of the trisodium salt of 15-isothiocyanato-4,7,10-tris(carboxymethyl)-2,17:12,14-dietheno-1,4,7,10,13-benzopentaazacyclopentadecine prepared in Preparation 8 and 200 parts by weight of methanol under argon at room temperature is rapidly added a solution of one equivalent of methylhydrazine dissolved in 10 parts of methanol. After one hour at room temperature, the solvent is removed by evaporation at reduced pressure, the residue is triturated with 10 parts of anhydrous oxygen-free ether, and the solid is isolated by filtration.


25

Preparation 10. Tetrasodium 2-Methyl-3-thio-4-{5-([6,6"-di-bis(carboxymethyl)aminomethyl]-2:2',6':2"-terpyridin-4'-yl)-2-methoxyphenyl]}-semicarbazide



To a freshly prepared and stirred mixture of one part by weight of the tetrasodium salt of the TMT-isothiocyanate prepared in Preparation 6 and 200 parts by weight of methanol under argon at room temperature is rapidly added a solution of one equivalent of methylhydrazine dissolved in 10 parts of methanol. After one hour at room temperature, the solvent is removed by evaporation at reduced pressure, the residue is triturated with 10 parts of anhydrous oxygen-free ether, and the solid is isolated by filtration.


Preparation 11. Tetrasodium 2-methyl-3-thio-4-((6,6'-bis(N',N'-dicarboxymethyl-N-methylhydrazino)-2:2',6:2'-terpyridine-4'-yl)-2-methoxyphenyl)}semicarbazide



5 To a freshly prepared and stirred mixture of one part by weight of the tetrasodium salt of the THT-isothiocyanate prepared in Preparation 3 and 200 parts by weight of methanol under argon at room temperature is rapidly added a solution of one equivalent of methylhydrazine dissolved in
 10 10 parts of methanol. After one hour at room temperature, the solvent is removed by evaporation at reduced pressure, the residue is triturated with 10 parts of anhydrous oxygen-free ether, and the solid is isolated by filtration.

15 Preparation 12. Tetrasodium 2-Methyl-3-thio-4-[2,9-di-bis(carboxymethyl)aminomethyl-1,10-phenanthrolin-5-yl]semicarbazide.

SUBSTITUTE SHEET



To a freshly prepared and stirred mixture of one part by weight of the tetrasodium salt of the PheMT-isothiocyanate prepared in Preparation 5 and 200 parts by weight of methanol under argon at room temperature is rapidly added a solution of one equivalent of methylhydrazine dissolved in 10 parts of methanol. After one hour at room temperature, the solvent is removed by evaporation at reduced pressure, the residue is triturated with 10 parts of anhydrous oxygen-free ether, and the solid is isolated by filtration.

Preparation of Conjugates of an Antibody
with TMT and with THT

Part A - Procedure to Attach Chelates to Antibodies

TMT or THT was attached to antibodies by oxidation of the carbohydrate groups on the antibodies with NaIO_4 to produce aldehyde groups on the antibodies. A solution (110 mL) of 0.1 M NaIO_4 in purified water was added to one mL of a 4 mg/mL solution of B72.3* antibody in PBS (10 mM sodium phosphate, 0.15 M NaCl), pH 6.0. The pH of the resulting solution was readjusted to 6.0 and incubated for 1 hour at room temperature in the dark. After this incubation, excess periodate was removed by passing the antibody solution through a Sephadex G50 column that had been equilibrated with PBS, pH 6.0. One mL fractions were collected and the two fractions giving

the highest absorbance at 280 nm were pooled.

5 A 0.33 M. solution of each chelating agent (TMT or THT) was prepared by dissolving the solid chelating agent in PBS, pH 6.0. This raised the pH of the solution to over pH 9.0 so the pH was adjusted to about 6.8 with 6 M HCl. Two hundred μ L of each chelator solution was added to separate 2 mL portions of the antibody solution and the pH of each resulting solution was 10 adjusted to pH 6.0. These mixtures were incubated for 5 h at room temperature in the dark. Then NaCNBH₃ (Aldrich 29,694-5; 5 M in about 1 M NaOH) was added to each solution to a final concentration of 10 mM and the pH was adjusted to 6.0 to reduce the Schiff base formed by the aldehyde. The mixtures were incubated overnight 15 at room temperature. The solutions were then centrifuged (Eppendorf Model 5412) to remove undissolved chelate that precipitated overnight and were concentrated to 1 mL in a Centricon 30 (Amicon) 20 microconcentrator. These solutions were passed through Sephadex G50 columns to remove excess chelator and NaCNBH₃. One mL fractions were collected and the two fractions of each chelator solution containing the highest absorbance at 280 nm were pooled. The pooled 25 fractions were dialyzed against 0.01 M sodium acetate, 0.15 M NaCl, pH 6.0 with two changes of buffer. If further precipitates were formed during analysis, they were removed by centrifugation.

Typically, the conjugates produced by this 30 procedure had a ratio of TMT or THT to antibody of about 1.7.

35 * B72.3 is a well known antibody to colorectal tumor-associated antigens which was first disclosed by the National Institute of Health (NIH).

Part B - Analysis of Ratio of Chelator to Antibody

The concentrations of antibody in the conjugate solutions were determined using the BioRad protein assay using bovine immunoglobulin as the protein standard. The concentration of THT was determined spectrophotometrically using the measured extinction coefficient at 350 nm. For TMT-B72.3 conjugate a portion (about 0.5 mg) of the conjugate was put into 1 mL of 0.01 M sodium acetate, 0.15 M NaCl buffer, pH 6.0 and an excess of Eu^{+3} was added. The absorbance of this solution at 330 nm was used to determine the amount of chelate in these solutions.

Part C - In Vitro Functional Test (Immunocompetency Assay) of Antibody-Chelate Conjugates

The wells of Linbro EIA II Plus microtiter plates were coated with antigen by adding 100 μL /well of a solution of 4 $\mu\text{g}/\text{mL}$ of bovine submaxillary mucin (Sigma M4503), an antigen to the B72.3 antibody, in PBS and incubated for 1 h at room temperature. After washing the plate three times with PBS, pH 6.8, the wells were blocked by adding 200 μL /well of a 1% BSA (bovine serum albumin, Sigma A-7906)-PBS, pH 6.8 solution and incubated for 1 h at room temperature. The plates were again washed three times with PBS, pH 6.8. Samples of the conjugates were prepared at a concentration of 1×10^{-7} M in PBS, pH 6.8 containing 1% horse serum (Sigma S-7390), and dilutions in this same buffer were made from these solutions. One hundred mL of each sample dilution were added to wells, in duplicate. After a 1 h room temperature incubation, the plates were washed three times with PBS, pH 6.8 and then scored by the addition of 100 μL per well of a 1:1000 dilution of a rabbit anti-mouse F(ab')_2 -HRP conjugate (Jackson Labs, 315-035-047) in PBS, pH 6.8, containing 1% horse serum. After a 1 h room temperature incubation, the wells were washed three

times with PBS, pH 6.8. Color developed upon the addition of 100 μ L per well of ABTS-HRP substrate (Kirkegaard and Perry Labs, Product #506502 and 506402). The reaction was stopped by the addition of an equal volume of 4N H_2SO_4 . The color was read using a 414 nm filter in a Titertek Multiscan instrument after 15 min.

When tested by this procedure, the immunoconjugates of B72.3 with TMT or THT were found to have immunoreactivity comparable to native B72.3. The data are presented in Figures 1 and 2. Figure 1 also contains the curve for a scandium complex of the B72.3-THT conjugate prepared in the same manner as the Europium complex described hereinbefore.

Examples 1-2 Preparation and Evaluation of Radionuclide (^{111}In and ^{90}Y) Chelate Complexes of a TMT Conjugate and, as a Comparative Control, of a DTPA Conjugate

In Vivo Functional Test of Chelate-Antibody Conjugates-Biodistribution Experiments

1. Test Materials. The immunoconjugates tested were a B72.3-TMT conjugate prepared as described above or a B72.3-DTPA conjugate in which the DTPA was attached (through a linker arm) to oxidized carbohydrate on the antibody using procedures similar to those described for TMT attachment.

2. Labeling Conjugates with Radioisotopes. The antibody-chelate conjugates to be labeled with either ^{111}In or ^{90}Y were in phosphate buffered saline, pH 6.0. The radionuclides (about 1 mCi) were added to the conjugate solution (1 mL containing about 1 mg of antibody-chelate conjugate), mixed, and incubated about 30 min at room temperature. Any non-chelated metal was removed from each conjugate preparation by HPLC gel filtration (TSK3000SW column). The column effluent was monitored for both protein (OD280 nm) and radioactivity

(radioactivity monitor). The specific activity of the purified labeled conjugates was calculated from this data. The conjugates were used for biodistribution studies immediately.

5 3. General Procedure. Five mice were used in each test group for the ^{90}Y biodistribution study and 3 mice in each group for the ^{111}In study. Target doses for each mouse were 10 μg of radioactive conjugate per dose, 20-100 mg antibody per μL , greater
10 than 50 μCi per dose. Both non-tumor bearing and tumor bearing mice were injected on Day 0.

 4. Tumor Growth Initiation. Each nude mouse (nu/nu: Swiss Background, Taconic Farms, Germantown, NY) was injected subcutaneously in the left rear flank
15 with one million LS174T cells in exponential growth phase, in about 0.2 mL of sterile medium or saline from cell culture. The mice were examined for tumor growth until tumors became measurable. Thereafter, tumors were measured to the nearest 0.1 mm, across two
20 perpendicular diameters (length x width) using digital calipers until the product of the length times the width was between 50 and 100 square millimeters for all the mice.

 5. Injection Protocol. On a desirable day following tumor cell inoculation (day 0), each mouse
25 was injected with the labeled conjugate. The radiolabeled antibody for each dose was drawn up into a separate 1 cc insulin syringe with a 28G, 1/2 inch needle for administration. Three 10 μL aliquots of
30 each test material were saved for gamma counting to determine the injected dose.

 Each syringe containing test material was numbered in order of administration. Each syringe was weighed and counted in a dose calibrator set to quantitate the
35 radionuclide. This information was recorded. The mice were anesthetized by injecting i.p. 0.15 mL of a

sterile solution containing ketamine HCl at 11 mg/mL and xylazine at 3 mg/mL. The test material was injected via the retroorbital venous sinus. Each mouse was counted in the dose calibrator immediately after injection, and the syringes were reweighed, recounted and the data recorded.

6. Methods of Evaluation. Each animal was weighed to the nearest 0.1 g on Day 0 and just prior to dissection. Immediately prior to imaging and/or dissection, each animal was counted in the dose calibrator following standard operating procedures. Each carcass was counted after dissection. Immediately prior to dissection, each animal had its tumor measured to the nearest 0.1 mm across two perpendicular diameters (length x width) using digital calipers. A blood sample was collected from each animal into a tared culture tube and then the tube was reweighed. Each animal was sacrificed by cervical dislocation. Organs (lungs, spleen, liver, colon, right kidney, left kidney, tumor, bone marrow, muscle) were dissected and trimmed to remove extraneous tissue and then weighed to the nearest milligram. Radioactivity of the dissected organs was determined by gamma counting.

7. Results. The biodistributions shown in Figure 3 are for six mice four days after injection of the radioactive immunoreagents. The data from the ^{111}In biodistribution test indicate that the radiolabeled TMT conjugate of B72.3 did target the radioactivity to the tumor. Although liver and kidney levels were slightly greater than the B72.3-DTPA- ^{111}In conjugate, the B72.3-TMT- ^{111}In conjugate could be used for ^{111}In tumor targeting.

The biodistributions shown in Figure 4 are eight days after injection of the radioactive immunoreagent. The average value for each group of five mice is graphed for the tissue type examined. The data from

the ^{90}Y biodistribution show that the B72.3-TMT- ^{90}Y conjugate targeted ^{90}Y to the tumor as well as the B72.3-DTPA- ^{90}Y conjugate. However, considerably less ^{90}Y was found in the femur when the B72.3-TMT- ^{90}Y conjugate was used compared to the B72.3-DTPA- ^{90}Y conjugate. Since the femur (bone marrow) is the dose-limiting organ for ^{90}Y therapy, this result indicates that TMT is a better chelate for antibody targeted radionuclide therapy using this isotope. Of the other tissues examined, only the blood was found to have a higher ^{90}Y level when TMT was the chelate used to prepare the immunoconjugate. This is believed to be due to superior in vivo stability of the B72.3-TMT- ^{90}Y complex compared to the B72.3-DTPA- ^{90}Y complex.

Survival Study

If the B72.3-TMT- ^{90}Y complex decreases the amount of ^{90}Y that gets to the bone marrow compared to B72.3-DTPA- ^{90}Y and the bone marrow is the dose-limiting tissue, then mice inoculated with the conjugates should find the B72.3-TMT- ^{90}Y conjugate less toxic than those inoculated with B72.3-DTPA- ^{90}Y . The general procedures to test this hypothesis (survival study) were the same as those described for the biodistribution study with the following exceptions. There were 8 tumor bearing nude mice in each test group. The injection schedule was as follows: 200 μCi , 120 μCi , and 120 μCi were given on days 0, 4 and 8, respectively, for both the B72.3-TMT- ^{90}Y or B72.3-DTPA- ^{90}Y conjugate (low dose regimen). Two other sets of mice received 400 μCi , 320 μCi and 320 μCi of the two immunoconjugates on days 0, 4 and 8, respectively (high dose regimen). The survival of the mice was observed (no biodistributions of the test material were performed). Results (Figure 5) indicate that the survival of the mice was prolonged if ^{90}Y is administered using the B72.3-TMT conjugate.

TMT-Immunoconjugate with anti-tumor specificity


5 TMT or a suitable derivative thereof can be
conjugated to an antibody molecule to yield an
antibody-TMT conjugate molecule that displays the
ability to bind to a target antigen recognized by the
antibody variable region. Such a conjugate molecule
10 can be used to deliver a radioisotope that is chelated
by the TMT moiety in order to localize and/or treat the
lesion that is targeted by such an immunoconjugate. In
one preferred embodiment, the antibody is selected such
that it has a broad reactivity with an antigen molecule
15 expressed on tumor cells, thereby providing an
antibody-TMT conjugate that can deliver radioisotope to
the tumors for therapeutic or diagnostic purposes.
ING-1 is a chimeric antibody (described in
International Patent Publication Number WO 90/02569,
20 dated 22 March 1990) consisting of a murine variable
region and a human immunoglobulin constant region. The
antibody molecule is produced by culturing a mouse
myeloma cell line expressing the chimeric antibody
essentially as described in the above-referenced
publication. The chimeric ING-1 antibody is used at a
25 concentration of 5.2 mg/ml in 50mM sodium acetate
buffer at pH 5.6 and supplemented with 150mM NaCl. To
1.15 ml of antibody solution is added a solution of
50mM sodium borate at pH 9.0 supplemented with 100mM
sodium chloride to a final total volume of 2.5 ml. The
30 solution is applied to a PD-10 chromatography column
equilibrated with the sodium borate buffer added to the
antibody solution. The antibody is eluted off the
column with 3.5 ml of the same sodium borate buffer.
The eluate is concentrated using a Centricon-30 to a
35 concentration of approximately 4.0 mg ING-1 chimeric
antibody per milliliter solution. The solution of the

NCS derivative of TMT (Preparation 6), i.e., TMT wherein the NH_2 is replaced with NCS, (designated herein as TMT-NCS) is prepared in the sodium borate buffer at a concentration of 10mg/ml and added to the antibody solution to a final concentration of 138 μM of TMT-NCS. The solution is gently mixed and incubated in the dark at ambient temperature (approx. 22°C) overnight (approx. 12 hr). The ING-1:TMT conjugate is separated from free TMT-NCS and other low molecular weight products using a Superose 12 HPLC column equilibrated and eluted in 50mM sodium acetate buffer at pH 5.6 supplemented with 150mM sodium chloride. The TMT immunoconjugate is then tested for its ability to bind to the tumor cell antigen and also to bind Yttrium-90 isotope to demonstrate that the conjugate can be radiolabeled in the TMT moiety and can target the tumor cell.

The invention has been described in detail with particular reference to certain preferred embodiments thereof, but it will be understood that variations and modifications can be effected within the spirit and scope of the invention.

What is claimed is:

1. A targeting radioactive immunoreagent comprising
 - a metal radionuclide ion,
 - a complexing agent, and
 - an immunoreactive group covalently bonded to said complexing agent,
 said complexing agent having the structure



wherein

R represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group;

R¹ represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group;

R² represents hydroxy, carboxy, hydroxyalkyl,

carbonyliminodiacetic acid, methyleneiminodiacetic acid, methylenethioethyleneiminodiacetic acid, hydrozinylylidenediacetic acid, or a salt of such acids, or the two R^2 groups, taken together, represent the atoms necessary to complete a macrocyclic ring structure containing at least one heteroatom coordinating site and at least one alkylene group forming part of the ring structure;

R^3 represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group;

R^4 represents hydrogen or a protein reactive group;


n is 0 to 4;

o is 0 or 1;

m is 0 or 1;


provided that at least one of n and m is 0 and at least one of R , R^1 , R^3 and R^4 is a protein reactive group.

2. The immunoreagent of claim 1 comprising a complexing agent having the structure:



wherein R^1 , R^2 and R^3 are as defined in claim 1.


3. The immunoreagent of claim 1 comprising a complexing agent having the structure:



wherein R, R¹, R² and R³ are as defined in claim 1 and n is 0, 1, 2, 3 or 4.

4. The immunoreagent of claim 1 wherein said complexing agent is TMT or THT.

5. The immunoreagent of claim 1 comprising a complexing agent having the structure:



wherein R², R³ and R⁴ are as defined in claim 1.

6. The immunoreagent of claim 1 wherein said protein reactive group is selected from the group consisting of amino, alkylamino, arylamino, hydrazino, alkylhydrazino, arylhydrazino, carbazido, semicarbazido, thiocarbazido, thiosemicarbazido, sulfhydryl, sulfhydrylalkyl, sulfhydrylaryl, hydroxy, carboxy,

carboxyalkyl, carboxyaryl, active halogen containing groups, 2-leaving group-substituted ethylcarbonyl, vinylsulfonyl, vinylcarbonyl, epoxy, isocyanato, isothiocyanato, aldehyde, aziridine, succinimidoxycarbonyl; activated acyl groups, and mixed anhydrides.

7. The immunoreagent of claim 1, wherein said immunoreactive group is selected from the group consisting of enzymes, amino acids, polypeptides, proteins, lipoproteins, glycoproteins, hormones, drugs, steroids, vitamins, polysaccharides, viruses, protozoa, fungi, parasites, rickettsia, molds, blood components, tissue and organ components, pharmaceuticals, haptens, lectins, toxins, nucleic acids, antibodies, antigenic materials, avidin and biotin.

8. The immunoreagent of claim 1, wherein said immunoreactive group is an antibody.

9. The immunoreagent of claim 1 wherein said immunoreactive group is an antibody selected from the group consisting of B72.3, 9.2.27, D612, UJ13A, NRLU-10, 7E11C5, CC49, TNT, PR1A3, ING-1, B174 and B43 antibodies.

10. The immunoreagent of claim 9 wherein said antibody is ING-1.

11. The immunoreagent of claim 1 wherein said metal radionuclide ion is $^{90}\text{Y}^{+++}$.

12. A diagnostic imaging composition comprising the radioactive immunoreagent of claim 1.

13. A therapeutic composition comprising the radioactive immunoreagent of claim 1 and a pharmaceutically acceptable carrier therefor.


14. A method for diagnostic imaging a site in a patient comprising the steps of

a) administering to the patient an effective amount of the radioactive immunoreagent of claim 1 capable of targeting said site, and

b) imagewise activating a radiation sensitive element or device with the radiation emitted from the targeted site.

15. A method for treating disease sites in a patient comprising the steps of administering to the patient or a specimen from the patient an effective amount of a therapeutic composition comprising the radioactive immunoreagent of claim 1 and a pharmaceutically acceptable carrier therefor.

16. A composition of matter having the structure



wherein

R^1 represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group;

R^2 represents hydroxy, carboxy, hydroxyalkyl, carbonyliminodiacetic acid, methyleneiminodiacetic acid, methylenethioethyleneiminodiacetic acid, hydrozinylylidenediacyetic acid, or a salt of such acids, or the two R^2 groups, taken together, represent the atoms necessary to complete a macrocyclic ring structure containing at least one heteroatom coordinating site and at least one alkylene group forming part of the ring structure;


R^3 represents hydrogen, alkyl, alkoxy, alkylthio,

alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl, or a protein reactive group;


R⁵ represents alkyl or alkoxy; and

R⁶ represents a protein reactive group.

17. The composition of claim 16 having the structure



18. A composition of matter having the structure



wherein

R² represents hydroxy, carboxy, hydroxyalkyl, carbonyliminodiacetic acid, methyleneminodiacetic acid, methylenethioethyleneiminodiacetic acid,

hydrozinylylidenediacetic acid, or a salt of such acids, or the two R^2 groups, taken together, represent the atoms necessary to complete a macrocyclic ring structure containing at least one heteroatom coordinating site and at least one alkylene group forming part of the ring structure; and

R^3 represents hydrogen, alkyl, alkoxy, alkylthio, alkylamino, alkylformamido, aryl, aryloxy, heterocyclyl or a protein reactive group; and

R^4 represents hydrogen or a protein reactive group, provided that at least one of the R^4 groups is a protein reactive group.

FIG. 1

IMMUNOCOMPETENCY, BSM (MUCIN) PLATES




FIG. 2




FIG. 3
LS174T TUMOR BEARING ANIMALS




FIG. 4

BIODISTRIBUTION OF Y-90 IN NUDE MICE
LS174T CELL XENOGRAFT




FIG. 5
SURVIVAL CURVE




FIG. 6




FIG. 7

